100

But, aside from the interest attached to this new genus of the Volvociniae from the botanical point of view, it may be found to have important relations to odors and tastes in water supplies, when it will become the enemy of engineers and water commissioners, as other members of this group have done before. For example, Volvox globator has caused much trouble in Rochester, N. Y., by imparting a disagreeable fishy odor to the city water supply, and in Massachusetts Pandorina and Eudorina have caused similar troubles on a smaller scale. Plodorina, coming as it does between Volvox and Eudorina in the classification, may be looked upon with suspicion in this respect, if it ever infects a water supply in a sufficient quantity. On account of the filthy condition of the canal water in which it was found in Lafayette, and the number of other forms growing with it, no idea could be formed as to the nature of the odor, if any, of Pleodorina.

FORMS OF XANTHIUM CANADENSE AND X. STRUMARIUM. By J. C. ARTHUR, In the absence of the author the outline of the paper was presented by Mr. Wm. Stuart and photographs of the two species were shown. The species in their most typical forms differ widely in the outline of the leaf and character and size of the burs. X. Canadense has a flowing sub-entire outline to the leaf, and large, strongly hispid fruit covered thickly with prickles, while X. strumarium has dentate leaves and smaller glabrons fruit with fewer prickles. All gradations exist between the two types, due possibly to hybridization.

NOTES ON WOOD SHRINKAGE. BY M. J. GOLDEN.

The increase or diminution in size of a piece of wood, due to its possession of a greater or less amount of moisture, is well known, as is also the fact that this change in size may be accompanied by the expenditure of a great deal of force. If an unseasoned piece of wood has two sides fastened rigidly so that it can not shrink across the grain, and then be exposed to a current of comparatively dry air, it will very soon break, the break being in the direction of the length of the cells of which the wood fibers are composed; or if a piece of dry wood be confined rigidly to prevent any *increase* in size and then be saturated with moisture, it will tend to swell and the force will be sufficient to crush the fibers where they are in contact with whatever confines them.