

Higher Fungi of Marion County, Indiana

JOHN O. COTTINGHAM, Indianapolis, Indiana

The following is a list of fleshy, leathery and woody fungi which have been found by my assistants and myself since my report to the Indiana Academy of Science, October 1947. Our report at that time listed 192 species. We are now adding 46 more or a total of 238 species thus far found and identified in Marion County. Many species that should be indigenous have not yet been found and while this list is very incomplete, it does give the mycologist a good idea of what he may expect to find most frequently in the center of Indiana.

AGARICACEAE

Agaricus arvensis, Fr.; *Collybia fusipes* (Bull); *Collybia maculata*, A. & S.; *Collybia hariolorum*, Fr.; *Clitopilus popinalis*, Fr.; *Clitocybe dealbata*, Fr.; *Cortinarius duracinus*, Fr.; *Hygrophorus flavo-discus*, Frost.; *Lactarius piperatus*, Fr.; *Lentinus Ursinus*, Fr.; *Lenzites vialis*, Pk.; *Mycena alcalina*, Fr.; *Marasmius foetidus*, Sow.; *Panus stipticus*, Bull.; *Pholiota rugosa*, Pk.; *Pholiota squarrosa*, Mull.; *Pholiota mutabilis*, Schaeff.; *Pholiota curvipes*, Fr.; *Psilocybe foenicicii*, Fr.; *Russula ochrophylla*, Pk.; *Tricholoma sulphureum*, Fr.; *Tricholoma album*, Schaeff.; *Collybia hygrophoroides*, Pk.

POLYPORACEAE

Polyporus Berkeleyi, Fr.; *Polyporus pensylvanicus*, Sumst.; *Polyporus chioneus*, Fr.; *Polyporus radiatus*, Fr.; *Polyporus pergamenus*, Fr.; *Polyporus frondosus*, Fr.; *Polyporus fumosus*, Fr.; *Polyporus tephroleucis*, Fr.; *Polyporus sanguineus*, Fr.; *Boletus bicolor*, Pk.; *Boletus luteus*, L.; *Poria corticola*, Fr.; *Poria attenuata*, Pk.; *Poria medulapanis*, Pers.; *Trametes rigida*, Bert. and Mont.

LYCOPERDACEAE—*Calvatia gigantea*, Pers.

HYDNACEAE

Hydnum erinaceus, Bull.; *Hydnum albonigrum*, Pk.; *Hydnum ochraceum*, Pers.

THELOPHORACEAE—*Sterum rameale*, Schw.

ASCOMYCETES—*Peziza vesiculosa*, Bull.